No.

200500349

# <u>THER UNITED STATES OF AMERICA</u>

TO ALL TO WHOM THUSE: PRESENTS: SHATE COME:

**Tonisiana State Unibersity Agricultural Center** 

MICCENS, THERE HAS BEEN PRESENTED TO THE

## Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLEMSHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE SIGHT TO EXCLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR FOREIGN IT, OR EXPORTING IT, OR CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE VERLY PURPOSE. OR CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE ABOVE SE, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT VARIETY THEREFROM, TO THE EXTENT BY THE PLANT VARIETY PROTECTION ACT. IN THE UNITED STATES SEED OF THIS VARIETY (I) SOLD BY VARIETY NAME ONLY AS A CLASS OF CERTIFIED SEED AND (2) SHALL CONFORM TO THE NUMB.

**RICE** 

'CL131'

In Jestimonn Marrot, I have hereunto set my hand and caused the seal of the Hant Bariety Protection Office to be affixed at the City of Washington, D.C. this fifth day of June, in the year two thousand and six.

Attest:

ET SEQ

Removed Commissioner

Commissioner Plant Variety Protection Office Agricultural Marketing Service Agriculture

(See reverse for instructions and information collection burden statement)

#### INSTRUCTIONS

GENERAL: To be effectively filed with the Plant Variety Protection Office (PVPO), ALL of the following item's must be received in the PVPO? (1) Completed application form signed by the owner; (2) completed exhibits A, B, C, E; (3) for a seed reproduced variety at least 2,500 viable untreated seeds, for a hybrid variety at least 2,500 untreated seeds of each line necessary to reproduce the variety, or for tuber reproduced varieties verification that a viable (in the sense that it will reproduce an entire plant) tissue culture will be deposited and maintained in an approved public repository; (4) check drawn on a U.S. bank for \$3,652 (\$432 filing fee and \$3,220 examination fee), payable to "Treasurer of the United States" (See Section 97.6 of the Regulations and Rules of Practice.) Partial applications will be held in the PVPO for not more than 90 days, then returned to the applicant as unfiled. Mail application and other requirements to Plant Variety Protection Office, AMS, USDA, Room 401, NAL Building, 10301 Baltimore Avenue, Beltsville, MD 20705-2351. Retain one copy for your files. All items on the face of the application are self explanatory unless noted below. Corrections on the application form and exhibits must be initiated and dated. DO NOT use masking materials to make corrections. If a certificate is allowed, you will be requested to send a check payable to "Treasurer of the United States" in the amount of \$432 for issuance of the certificate. Certificates will be issued to owner, not licensee or agent.

Plant Variety Protection Office Telephone: (301) 504-5518 FAX: (301) 504-5291

200500349

Homepage: http://www.ams.usda.gov/science/pvpo/pvp.htm

To avoid conflict with other variety names in use, the applicant must check the appropriate recognized authority and provide evidence that name has been cleared by the appropriate recognized authority before the Certificate of Protection is issued. For example, for agricultural and vegetable crops, contact: Seed Branch, AMS, USDA, 10301 Baltimore Avenue, Suite 401 NAL Building, Beltsville, MD 20705. Telephone: (301) 504-5682 http://www.ams.usda.gov/isg/seed.htm.

#### ITEM

- 19a. Give:
- (1) the genealogy, including public and commercial varieties, lines, or clones used, and the breeding method;
- (2) the details of subsequent stages of selection and multiplication;
- (3) evidence of uniformity and stability; and
- (4) the type and frequency of variants during reproduction and multiplication and state how these variants may be identified
- 19b. Give a summary of the variety's distinctness. Clearly state how this application variety may be distinguished from all other varieties in the same crop. If the new variety is most similar to one variety or a group of related varieties:
  - (1) identify these varieties and state all differences objectively;
  - (2) attach statistical data for characters expressed numerically and demonstrate that these are clear differences; and
  - (3) submit, if helpful, seed and plant specimens or photographs (prints) of seed and plant comparisons which clearly indicate distinctness.
- 19c. Exhibit C forms are available from the PVPO Office for most crops; specify crop kind. Fill in Exhibit C (Objective Description of Variety) form as completely as possible to describe your variety.
- 19d. Optional additional characteristics and/or photographs. Describe any additional characteristics that cannot be accurately conveyed in Exhibit C. Use comparative varieties as is necessary to reveal more accurately the characteristics that are difficult to describe, such as plant habit, plant color, disease resistance, etc.
- 19e. Section 52(5) of the Act requires applicants to furnish a statement of the basis of the applicant's ownership. An Exhibit E form is available from the PVPO.
- 20. If "Yes" is specified (seed of this variety be sold by variety name only, as a class of certified seed), the applicant MAY NOT reverse this affirmative decision after the variety has been sold and so labeled, the decision published, or the certificate issued. However, if "No" has been specified, the applicant may change the choice. (See Regulations and Rules of Practice, Section 97.103).
- 23. See Sections 41, 42, and 43 of the Act and Section 97.5 of the regulations for eligibility requirements.
- 24. See Section 55 of the Act for instructions on claiming the benefit of an earlier filing date.
- 22. CONTINUED FROM FRONT (Please provide a statement as to the limitation and sequence of generations that may be certified.)

The following generations may be certified: Breeder-Foundation-Registered-Certified

23. CONTINUED FROM FRONT (Please provide the date of first sale, disposition, transfer, or use for each country and the circumstances, if the variety (including any harvested material) or a hybrid produced from this variety has been sold, disposed of, transferred, or used in the U.S. or other countries.)

Sale - Foundation and Registered - March 1, 2005, USA -- transferred to Uruguay July 2005

24. CONTINUED FROM FRONT (Please give the country, date of filing or issuance, and assigned reference number, if the variety or any component of the variety is protected by intellectual property right (Plant Breeder's Right or Patent).)

US Patent 6,943,280 Resistance to acetohydroxycid synthase-inhibiting herbicides/

Utility patent serial number 60/715,690 Filed September 9, 2005

NOTES: It is the responsibility of the applicant/owner to keep the PVPO informed of any changes of address or change of ownership or assignment or owner's representative during the life of the application/certificate. The fees for filing a change of address; owner's representative; ownership or assignment; or any modification of owner's name is specified in Section 97.175 of the regulations. (See Section 101 of the Act, and Sections 97.130, 97.131, 97.175(h) of the Regulations and Rules of Practice.)

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information collection is 0581-0055. The time required to complete this information collection is estimated to average 1.4 hours per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, sexual orientation, marital or family status, political beliefs, parental status, or protected genetic information. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whilten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964 (voice and TDD). USDA is an equal opportunity provider and employer.

 $\frac{\partial}{\partial x}$ 

## **EXHIBIT A**

## Origin and Breeding History

|      | CL131 – Development History<br>Pedigree: CL161//Kaybonnet/LA2031 |                               |  |  |  |  |
|------|--|-------------------------------|--|--|--|--|
| Year | Generation   | ID                            |  |  |  |  |
| 2000 | Cross F <sub>0</sub>   | 00CR387                       |  |  |  |  |
| 2001 | $F_1$  | 01T034                        |  |  |  |  |
| 2001 | F <sub>2</sub>   | 01B268-300 (Puerto Rico)      |  |  |  |  |
| 2002 | F <sub>3</sub>   | 02P2056 (Puerto Rico)         |  |  |  |  |
| 2002 | F <sub>4</sub>   | 0283305                       |  |  |  |  |
| 2002 | F <sub>5</sub>   | 02PR1351 (Puerto Rico)        |  |  |  |  |
| 2003 | F <sub>6</sub>   | CL002 (Clearfield Yield Test) |  |  |  |  |
| 2004 |  |                               |  |  |  |  |
| 2004 | - F <sub>8</sub>   | Foundation Seed Production    |  |  |  |  |

Details of stages of selection and multiplication: CL131 was developed from a modified program of single seed descent. From the original cross made in 2000, four plants were grown in 2001. The seed from these plants were bulked and used to grow a large  $F_2$  population at the Puerto Rico winter nursery (planted in August 2001). One panicle was selected from each of 150  $F_2$  plants and these were grown as  $F_3$  panicle rows in Puerto Rico (planted in December 2001). Row number 02P2056 was selected for advancement. Ten panicles were selected from this row and used to plant five  $F_4$  panicle rows on the Rice Research Station in May 2002. Five panicles were selected from row 0283305 and used to plant five  $F_5$  panicle rows in Puerto Rico in October 2002. Row 02PR1351 was selected for advancement and five panicles were selected from this row then the remaining seed was bulked. The bulked seed was used for testing and the five panicles were seeded for increase and purification purposes of the Rice Research Station in the summer of 2003. Seed was harvested from the five panicle rows and used to plant a seed increase in Puerto Rico in January 2004. This seed was reselected, harvested, and returned to the Rice Research Station. It was used to plant a Breeder/Foundation field on the Station in early June 2004. This seed was the origin of CL131 that has been provided to the industry.

CL131 has been observed for four generations of increase and multiplication and has exhibited a high level of uniformity and stability.

CL131 was originally selected in the F<sub>5</sub> generation as a semidwarf, very early imazethapyr-resistant line that displayed good yield potential and quality characteristics.

In each generation of multiplication and purification ( $F_5 - F_8$ ), the line was selected for uniformity and purity.

Variants observed and removed from increase fields of CL131 included any combination of the following: taller, earlier, later, medium grain, and intermediate grain. The total number of variants numbered less than 1 per 5,000 plants.

## Exhibit B

### **Statement of Distinctness**

CL131 is a very early maturing, imazethapyr-resistant long-grain rice variety. The line possesses the CFX18 gene for resistance to imazethapyr, which is the same gene found in CL161. CL131 was developed from the cross CL161//Kaybonnet/LA2031 made at the Rice Research Station in 2000 (00CR387). In testing, the line has averaged 91 cm in plant height compared with 104 for CL161.

CL131 most closely resembles the rice variety CL161. CL131 has an erect flag leaf angle at maturity while CL161 displays intermediate erectness at this growth stage. CL161 also averages 13 cm taller (at maturity) than CL131.

CL131 differs from all conventional rice varieties by the fact that it is resistant to imazethapyr and conventional varieties are not. It differs from CL121, CL141, and CFX51 in that it contains the CFX18 gene for imazethapyr resistance, which impacts a much higher level of resistance to imazethapyr than that found in the earlier Clearfield varieties.

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information collection is 0581-0055. The time required to complete this information collection is estimated to average 1.4 hours per response, including the time for reviewing instructions, searching existing data sources, gethering and maintaining the data needed, and complete the information collection of information collection is estimated. searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, sexual orientation, marital or family status, political beliefs, parental status, or protected genetic information. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964 (voice and TDD). USDA is an equal opportunity provider and employer.

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE SCIENCE AND TECHNOLOGY PLANT VARIETY PROTECTION OFFICE BELTSVILLE, MD 20705

Exhibit C

**OBJECTIVE DESCRIPTION OF VARIETY** Rice (Orvza sativa)

| NAME OF APPLICANT (S)  | 4   | EXPERIMENTAL DESIGNATION                    | VARIETY NAME          |       |
|--|---|---|-----------------------|-------|
| Louisiana State Universi  Agricultural Center  | CL020   |   | CL13                  | 31    |
| ADDRESS (Street and No. or RD No., City, State, and Zip  |   |   | FOR OFFICIAL USE O    |       |
| Rice Research Station  | cour, course,   |   | PVPO NUMBER           |       |
| 1373 Caffey Road   |   |   |                       |       |
| Rayne, LA 70578  |   |   | 2005                  | 00349 |
| PLEASE READ ALL INSTRUCTIONS C   | ADFEITI V.  |   |                       |       |
| Place the appropriate number that descrit<br>to descriptors developed by IBGR-IRRI R<br>readily by describing as many characters | bes the character of this variet<br>lice Advisory Committee and t |   |                       |       |
| 1. MATURITY: Days to Heading (Seedli   | ng to 50% Heading)  |   |                       |       |
| A. South: (Location: Lo  | ouisiana  | ) at <b>165</b>                             | kg/ha (Nitrogen Rate) | ·     |
| 84.8Number of Days   |   |   | <u> </u>              |       |
|  | e Vadata CT 1616  |   |                       |       |
|  | k Variety: CL1616   |   |                       |       |
| Days Same As Check   | k Variety:  | <del></del>                                 |                       |       |
| Days Later Than Check  | k Variety:  |   |                       |       |
| Maturity Class   | ery Early (85 Days or Less)<br>termediate (101 - 115)             | 2 = Early (86 - 100)<br>4 = Late (More Than | 115)                  |       |
| B. California: (Location:  |   | ) at  | kg/ha (Nitrogen Rate) |       |
| Number of Days   |   |   |                       |       |
| Days Earlier Than Check  | « Variety:  |   |                       |       |
|  | « Variety:  |   |                       |       |
| Days Later Than Check  | v Variety:  |   |                       |       |
|  | ery Early (90 Days or Less)<br>termediate (98 - 104)              | 2 = Early (91 – 97)<br>4 = Late (More Than  | 104)                  |       |
| 2. CULM:   |   |   |                       |       |
|  |   | est on the ground)                          | pen (About 60°)       |       |

|        |                                |   |   |   |                                      |   |          | EXHIBIT O (IC |
|--------|--------------------------------|---|---|---|--------------------------------------|---|----------|---------------|
| 2.     | CULM: (continued)              |   |   |   |                                      | <b>&amp;V U D V</b>                           | V J 4 9  |               |
|        | LENGTH                         |   |   |   |                                      |   |          | ÷             |
|        | 90 • 0 cm (Soil level          | to top of extended pa                     | nicle on main stem)   |   |                                      |   | •        |               |
|        | _13 • 0 cm Shorter Th          | an Check V                                | ariety: <u>CL161</u>  |   |                                      |   |          |               |
|        | Length Sa                      | me as Check V                             | ariety:   |   |                                      |   |          |               |
|        | • cm Longer                    | Than Check V                              | ariety:   |   |                                      |   |          |               |
|        | _1_ Height Class:              | 1 = Semi dwarf                            | 2 = Short   | 3 = Medium                              | 4 = Tall                             |   |          |               |
|        | _2_Internode Color (A          | fter Flowering):                          | 1 = Green   | 2 = Light Gold                          | 3 = Purple Lines                     | 4 = Purple                                    |          |               |
|        | _1_ Strength (Lodging          | Resistance):                              | 1 = Strong (no Lo<br>5 = Intermediate e<br>9 = Very Weak (A | (Most Plants Lodged                     |                                      | ely Strong (Most Plants I<br>ost Plants Flat) | .eaning) |               |
| 3.     | FLAG LEAF (After Head          | ding):                                    |   |   |                                      |   |          |               |
|        | <u>29</u> • <u>0</u> cm Length |   |   | <u>10</u> • <u>0</u> mm Width           |                                      |   |          |               |
|        | 1_Pubescence:                  | 1 = Glabrous                              | 2 = Intermediate  | 3 = Pubescent                           |                                      |   |          |               |
|        | 1_ Leaf Angle (After           | Heading):                                 | 1 = Erect   | 3 = Intermediate                        | 5 = Horizontal                       | 7 = Descending                                |          |               |
|        | _2_ Blade Color:               | 1 = Pale Green<br>5 = Purple Margins      | 2 = Green<br>6 = Purple Blotch                              | 3 = Dark Green<br>7 = Purple            | 4 = Purple Tips                      |   |          |               |
|        | _1_ Basal Leaf Sheath          | h Color:                                  | 1 = Green   | 2 = Purple Lines                        | 3 = Light Purple                     | 4 = Purple                                    |          |               |
| <br>4. | LIGULE:                        |   |   | *************************************** |                                      |   |          |               |
|        | 7 - 2 mm Longth (E             | rom hope of coller to                     | the tip of lete vegets                                      | dive etern)                             |                                      |   |          |               |
|        |                                |   |   |   |                                      |   |          |               |
|        | 1_ Color (Late Veget           |   | 1 = White   | 2 = Purple Lines                        | 3 = Purple                           |   |          |               |
|        | _1_Shape:                      | 1 = Acute to Acumi                        |   | 2 = 2-Cleft                             | 3 = Truncate                         |   |          |               |
|        | _1_ Collar Color (Late         | - • •                                     | 1 = Pale Green  | 2 = Green                               | 3 = Purple                           |   |          |               |
|        | 1_ Auricle Color (Late         | e Vegetative Stage):                      | 1 = Pale Green  | 2 = Purple                              |                                      |   |          |               |
| 5. I   | PANICLE:                       |   |   |   |                                      |   |          |               |
|        | <u>19</u> • <u>0</u> cm Length |   |   |   |                                      |   |          |               |
|        |                                | 1 = Compact 5                             | = Intermediate  | 9 = Open                                |                                      |   |          |               |
|        | 2 Secondary Branch             | aing: 1 = /                               | Absent  | 2 = Light                               | 3 = Heavy                            | 4 = Clustering                                |          | · .           |
|        | 2 Exsertion (Near M            | aturity): 1 = [                           | Less than 90%   | 2 = 90 - 99%                            | 3 = 100% Exserted                    | <b>1</b>                                      |          | •             |
|        | _1_ Axis:                      | 1 = Straight                              | 2 = Droopy  |   |                                      |   |          |               |
|        | 3 Shattering:                  | 1 = Very Low (Less<br>7 = Moderately High |   | 3 = Low (1 - 5%)<br>9 = High (More than |                                      | 25%)  |          |               |
|        | 3_ Threshability:              | 1 = Difficult 2                           | = Intermediate  | 3 = Easy                                |                                      |   |          |               |
| 3. (   | GRAIN (Spikelet):              |   | * - 40-do-do-   |   |                                      |   |          |               |
|        | _1_Awns (After Full He         | eading):                                  | 0 = Absent<br>7 = Long and Part                             | 1 = Short and l<br>ly Awned             | Partly Awned<br>9 = Long and Fully   | 5 = Short and Fully Av<br>Awned               | vned     |               |
|        | 6 Apiculus Color (At           | Maturity):                                | 1 = White<br>5 = Red Apex                                   | 2 = Straw<br>6 = Purple                 | 3 = Brown (Tawny)<br>7 = Purple Apex | 4 = Red                                       |          |               |
|        | _1_ Stigma Color:              | 1 = White 2                               | = Light Green   | 3 = Yellow 4                            | = Light Purple                       | 5 = Purple                                    | 6        |               |

|  |                            |   |                                    |                                 |                     |   |          | Exhibit C (Rice) |
|--|----------------------------|---|------------------------------------|---------------------------------|---------------------|---|----------|------------------|
| 6. GRAIN (Spikelet):   |                            |   |                                    |                                 |                     | 20050   | 0349     |                  |
| 0 Lemma and Palea Color (A   | t Maturity):               |   |                                    |                                 |                     |   |          |                  |
| 0 = Straw<br>3 = Brown Furrows on Stra<br>6 = Purple Spots on Straw<br>9 = Black | 4 = E $7 = F$              | Gold and/or G<br>Brown (Tawny<br>Purple Furrow<br>White |                                    | Straw Backgro                   | und                 | 2 = Brown Spots on<br>5 = Reddish to Ligh<br>8 = Purple |          |                  |
| Lemma and Palea Pubesce  |                            | Glabrous<br>Short Hairs                                 | 2 = Hairs on L<br>5 = Long Hairs   |                                 | 3 = Hairs           | s on Upper Portion                                      |          |                  |
| Spikelet Sterility (At Maturity  |                            | Highly Fertile<br>Highly Sterile                        | e (> 90%) 3 =<br>e (< 50% to Trace | Fertile (75 – 9<br>e) 9 = Cor   | 0%)<br>npletely Ste | 5 = Partly Sterile (5)<br>erile (0%)                    | ) – 74%) |                  |
| 7. GRAIN (Seed):   |                            |   |                                    |                                 |                     |   |          |                  |
| 2 Seed Coat (Bran) Color:  | 1 = White<br>5 = Red       | 2 ≔ Ligh<br>6 = Vari                                    | it Brown<br>able Purple            | 3 = Spe<br>7 = Pur              | ckled Brow<br>ple   | n 4 = Brown   | 1        |                  |
| 1_ Endosperm Type:   | 1 = Nonglutino             | ıs (Nonwaxy)  | 2 =                                | Glutinous (Wa                   | ıxy)                | 3 = Indeterminate                                       |          |                  |
| 1 Endosperm Translucency:  | 1 = Clear                  |   | 5 = Intermedia                     | te                              | 9 = Opac            | que   |          |                  |
| 1_ Endosperm Chalkiness:   | 0 = None<br>5 = Medium (10 | – 20% of Sa   | 1 = 9<br>mple) 9 = 1               | Small (Less th<br>arge (More th |                     |   |          |                  |
| _0_Scent (Aroma):  | 0 = Nonscented             | I   | 1 = Lightly Sce                    | nted                            | 2 = Scen            | ited  |          |                  |
| Shape Class (Length/Width I  | Ratio):                    |   |                                    |                                 |                     |   |          |                  |
| _3_ Paddy  | 1 = Short (2.2:1           | and Less)   | 2 = Medium (2                      | .3:1 to 3.3:1)                  | 3 = Long            | (3.4:1 and More)  |          | -                |
| 3 Brown  | 1 = Short (2.0:1           | and Less)   | 2 = Medium (2                      | .1:1 to 3.0:1)                  | 3 = Long            | (3.1:1 and More)  |          |                  |
| 3 Milled   | 1 = Short (1.9:1           | and Less)   | 2 = Medium (2                      | .0:1 to 2.9:1)                  | 3 = Long            | (3.0:1 and More)  |          |                  |
| Measurements:<br>Grain Form  | Length<br>(mm)             | Width<br>(mm)   | Thickness<br>(mm)                  | <b>.</b>                        | L/W<br>Ratio        | 1000 Grains<br>(grams)                                  |          |                  |
| Paddy  | 9.06                       | 2.53  | 2.03                               |                                 | 3.58                | 19.8  | •        |                  |
| Brown  | 7.14                       | 2.22  | 1.72                               |                                 | 3.22                | 17.1  |          |                  |
| Milled   | 6.62                       | 2.18  | 1.67                               |                                 | 3.04                | 16.8  |          |                  |
| 20 Milling Quality (% Hulls)   |                            | <u>67</u> Mi  | lling Yield (% Wi                  | ile Kernel (he                  | ad) Rice to         | Rough Rice)   |          |                  |
| <u>7.18</u> % Protein  |                            | <u>24.1</u> %   | 6 Amylose                          |                                 |                     |   |          |                  |
| Alkali Spreading Value:  | 1.5% KOH Sol               | ution   | 3.                                 | 91.7% K                         | OH Solution         | า   |          |                  |
| 5 Gelatination Temperature Ty  | pe:                        | 1 = High  | 5 = 1                              | ntermediate                     |                     | 7 = Low   |          |                  |
| Amylographic Paste Viscosity (Bra  | bender Units)              |   |                                    |                                 |                     |   |          |                  |
| Peak Hot Past  | е                          | Cooled F  | Paste 'E                           | Breakdown' 'S                   | etback'             |   |          |                  |
| <u>274.7</u> <u>163.3</u>  | _                          | 304.8   | <u> </u>                           | 111.3                           | <u></u>             |   |          |                  |
| 8. RESISTANCE TO LOW TEMPERTU  | JRE:                       |   | 2                                  |                                 |                     |   |          |                  |
| 2 Germination and Seedling Vi  | gor:                       | 1 = Low   | 2 = N                              | ledium                          | 3 = High            |   |          |                  |
| _2_ Flowering (Spikelet Fertility):  |                            | 1 = Low   | 2 = 1\                             | ledium                          | 3 = High            |   |          |                  |

9. SEEDLING VIGOR NOT RELATED TO LOW TEMPERATURE:

\_2\_ Vigor:

1 = Low

2 = Medium

3 = High

|              |           |                    |   |         |           |                                      |       |             |             | (See 4)              |                            | 8 8 8 8   | F 119 02        | Exhibit C (Ri |
|--------------|-----------|--------------------|---|---------|-----------|--------------------------------------|-------|-------------|-------------|----------------------|----------------------------|-----------|-----------------|---------------|
|              |           |                    |   |         |           | onal races for                       |       |             | -           |                      |                            |           |                 |               |
| 0 = lmr      | nune      | 1 :                | = Resistant                                 | 3 =     | Moderate  | ly Resistant                         | 5 = I | ntermediate | e 7=        | Moderate             | ly Suscept                 | tible     | 9 = Susceptible |               |
| Group        |           |                    | IB  |         |           | IC                                   |       | ID          |             | IE                   | IG                         | <u>IH</u> |                 |               |
| Number       | 1         | 5                  | 45  | 49      | 54        | 1                                    | 17    | 1           | 13          | 1                    | 1                          | 1         |                 |               |
| Resistance   |           |                    |   | 7       |           |                                      | 7     |             |             | <u>.</u>             |                            |           |                 |               |
| 11. RESIST   | ANCE T    | O OTHE             | R DISEASE                                   | <br>ES: |           |                                      |       |             |             |                      |                            | e.        |                 |               |
| 0 = lmn      | nune      | 1 :                | = Resistant                                 | 3 =     | Moderate  | ly Resistant                         | 5 = I | ntermediate | e 7=        | Moderate             | ly Suscept                 | tible     | 9 = Susceptible |               |
| _7_ N        | arrow Br  | own Lea            | f Spot (Cerc                                | spora o | ryzae)    |                                      |       | Agg         | regate Sh   | eath Spo             | (Rhizocto                  | nia Oryza | ae-sativae)     |               |
| _7_ Le       | eaf Smut  | (Entyloi           | ma oryzae)                                  |         |           |                                      |       | _7_ Stra    | ight Head   | ţ                    |                            |           |                 |               |
|              | rown Lea  |                    | Helminthosp<br>(=Bipolaris (<br>(=Drechsler | oryzae) |           |                                      |       | _7_ Keri    | nel Smut    |                      | ia horrida)<br>ı barclayar | na)       |                 | ·             |
| Le           | af Scald  | (Gerlac            | hia oryzae)                                 |         |           |                                      |       | Whit        | te Tip Ner  | natode (A            | phelencho                  | oides bes | seyi)           |               |
| Ho           | oja Blanc | a Virus            |   |         |           |                                      |       | Sten        | n Rot (Sc   | lerotium o           | ryzae)                     |           |                 |               |
| Sh           | eath Ro   | t (Saroci          | ladium oryza                                | re)     |           |                                      |       |             |             |                      |                            |           |                 |               |
| Py           | rthium Se | edling E           | Blight ( <i>Pythi</i> u                     | ım sp.) |           |                                      |       | Bact        | erial Bligh | nt (Xantho           | monas ca                   | mpestris  | pv. oryzae)     |               |
|              | neath Sp  | ot ( <i>Rhiz</i> o | octonia oryz                                | ae)     |           |                                      |       | 7 She       | ath Blight  | (Rhizocto            | nia solani                 | )         |                 |               |
| Ot           | her:      |                    |   |         |           |                                      |       |             |             |                      |                            |           |                 |               |
| 12. INSECT   | RESIST    | ANCE:              |   |         |           |                                      |       |             |             |                      |                            |           |                 |               |
| 0 = Imm      | une       | 1 =                | Resistant                                   | 3 =     | Moderatel | y Resistant                          | 5 = I | ntermediate | 7 =         | Moderate             | ly Suscept                 | ible      | 9 = Susceptible | 9             |
| _ <u></u> Gr | asshopp   | er                 |   |         |           |                                      |       | 9 Rice      | Stink Bu    | g (Oegalı            | ıs pugnax)                 | •         |                 |               |
| Ric          | ce Leafh  | opper              |   |         |           |                                      |       | Swa         | rm Caterp   | oillar               |                            |           |                 |               |
| Ric          | ce Hispa  |                    |   |         |           |                                      |       | 9 Rice      | Water W     | eevil ( <i>Lis</i> : | sorhoptrus                 | oryzoph   | ilus)           |               |
| Rice Midge   |           |                    |   |         |           | Rice Stalk Borer (Chilo plejadellus) |       |             |             |                      |                            |           |                 |               |

13. OTHER DESCRIPTORS: If there are other characters that describe this variety, please indicate below:

#### **REFERENCES**

\_\_\_ Sugarcane Borer (Diatraea saccharalis)

- C. R. Adair et al. 1972. Rice in the United States: Varieties and Production. USDA Handbook No. 289 (Rev.), 124 pp.
- J. G. Atkins et al. 1967. An International Set of Rice Varieties for Differentiating Race of Pyricularia Oryzae. Phytopath. 57:297-301.
- IBPGR-IRRI Rice Advisory Committee. 1980. Descriptors for Rice Oryzae sativa L. International Rice Research Institute. 21 pp.
- K. C. Ling and S. H. Ou, 1969. Standardization of the International Race Numbers of Pyricularia Oryzae. Phytopath. 59:339-342.
- B. D. Webb et al. 1985. Utilization Characteristics and Qualities of United States Rice. In Proceedings on Rice Grain Quality and Marketing. International Rice Research Institute (IRRI), Los Branos, Philippines. P. 25-35.

\_-\_ Least Skipper

Table 1. Average main crop yields (lb/A) for CL131 and CL161 across several trials at multiple locations in Louisiana (2003-2004).

| YEAR | THEST  | CP131 | CL:161 |
|------|--|-------|--------|
|      | CLRP - RRS                                   | 8830  | 8621   |
| 2003 |  |       |        |
|      | CL - LAKE ARTHUR                             | 5952  | 5461   |
|      | 2003 Average                                 | 7391  | 7041   |
|      | CL - RRS                                     | 6524  | 6511   |
|      |  |       |        |
|      | CL - ACADIA                                  | 6520  | 5993   |
|      | CL - EVANGELINE                              | 7172  | 7436   |
|      | CL - JEFF DAVIS                              | 4963  | 4974   |
|      | CL - LAKE ARTHUR                             | 6082  | 5875   |
|      | CL - PINE ISLAND                             | 4603  | 4777   |
| 2004 | CLPY - RRS                                   | 6804  | 7693   |
|      | CA - RRS                                     | 7206  | 7290   |
|      | CA - ACADIA                                  | 6712  | 6286   |
|      | CA - EVANGELINE                              | 7092  | 7149   |
|      | CA - LAKE ARTHUR                             | 5998  | 5762   |
|      | CA - MOREHOUSE                               | 8195  | 7847   |
|      | CA - PINE ISLAND                             | 5158  | 5152   |
|      | 2004 Average                                 | 6387  | 6365   |
|      | <u>*************************************</u> |       |        |
| •    | Grand Mean                                   | 6520  | 6455   |

Table 2. Whole rice yield (%) for CL131 and CL161 across several trials at multiple locations in Louisiana (2003-2004).

| YEAR | TEST             | CL131 | CL161 |
|------|------------------|-------|-------|
| 2003 | CLRP - RRS       | 69.7  | 68.9  |
|      | CL - LAKE ARTHUR | 66.9  | 63.5  |
|      | 2003 Average     | 68.3  | 66.2  |
|      |                  |       |       |
|      | CL - RRS         | 64.4  | 61.7  |
|      | CL - ACADIA      | 66.9  | 67.4  |
|      | CL - LAKE ARTHUR | 69.5  | 66.7  |
| 2004 | CLPY - RRS       | 66.4  | 69.1  |
|      | CA - RRS         | 64.6  | 62.9  |
|      | CA - ACADIA      | 67.2  | 67.4  |
|      | CA - LAKE ARTHUR | 68.8  | 67.9  |
|      | 2004 Average     | 66.8  | 66.2  |
|      |                  |       |       |
|      | Grand Mean       | 67.2  | 66.2  |

Table 3. Total rice yield (%) for CL131 and CL161 across several trials at multiple locations in Louisiana (2003-2004).

| -    |                  |          |       |
|------|------------------|----------|-------|
| YEAR | TEST             | CL131    | CL161 |
| 2003 | CLRP - RRS       | 73.1     | 72.4  |
|      | CL - LAKE ARTHUR | 73.6     | 70.5  |
|      | 2003 Average     | 73.4     | 71.5  |
|      |                  |          |       |
|      | CL - RRS         | 69.8     | 67.7  |
|      | CL - ACADIA      | 72.3     | 71.1  |
|      | CL - LAKE ARTHUR | 73.1     | 71.0  |
| 2004 | CLPY - RRS       | 71.4     | 72.2  |
|      | CA - RRS         | 69.9     | 69.0  |
|      | CA - ACADIA      | 72.2     | 71.1  |
|      | CA - LAKE ARTHUR | 73.6     | 72.5  |
|      | 2004 Average     | 71.8     | 70.7  |
| ٠,   |                  | <u> </u> | ·     |
|      | Grand Mean       | 72.1     | 70.8  |

Table 4. Mean plant height (cm) for CL131 and CL161 across several trials at multiple locations in Louisiana (2003-2004).

| YEAR | TEST             | CL131 | CLIGI |
|------|------------------|-------|-------|
| 2002 | CLRP - RRS       | 101   | 108   |
| 2005 | CL - LAKE ARTHUR | 95    | 108   |
|      | 2003 Average     | 98    | 108   |

| (    |                  |    |     |
|------|------------------|----|-----|
|      | CL - RRS         | 93 | 115 |
|      | CL - ACADIA      | 90 | 103 |
|      | CL - EVANGELINE  | 93 | 103 |
|      | CL - JEFF DAVIS  | 85 | 98  |
|      | CL - LAKE ARTHUR | 88 | 98  |
| 2004 | CL - PINE ISLAND | 88 | 98  |
| 2004 | CLPY - RRS       | 90 | 103 |
|      | CA - RRS         | 90 | 108 |
|      | CA - ACADIA      | 88 | 103 |
|      | CA - EVANGELINE  | 90 | 98  |
|      | CA - LAKE ARTHUR | 88 | 103 |
|      | CA - PINE ISLAND | 85 | 95  |
|      | 2004 Average     | 89 | 102 |

| Cuond Moon     | 00 | 403 |
|----------------|----|-----|
| │ Grand Mean │ | 90 | 103 |

Table 5. Mean number of days to 50% heading for CL131 and CL161 across several trials at multiple locations in Louisiana (2003-2004).

| YEAR | TEST             | CL131 | CL161 |
|------|------------------|-------|-------|
| 2003 | CLRP - RRS       | 77    | 77    |
|      | CL - LAKE ARTHUR | 80    | 79    |
|      | 2003 Average     | 79    | 78    |

|        |                  |    | ·  |
|--------|------------------|----|----|
|        | CL - RRS         | 89 | 95 |
|        | CL - ACADIA      | 88 | 93 |
|        | CL - EVANGELINE  | 88 | 89 |
| 12.000 | CL - JEFF DAVIS  | 89 | 92 |
|        | CL - LAKE ARTHUR | 88 | 90 |
|        | CL - PINE ISLAND | 83 | 84 |
| 2004   | CLPY - RRS       | 80 | 83 |
|        | CA - RRS         | 88 | 94 |
|        | CA - ACADIA      | 87 | 93 |
|        | CA - EVANGELINE  | 87 | 87 |
|        | CA - LAKE ARTHUR | 84 | 88 |
|        | CA - MOREHOUSE   | 83 | 85 |
|        | CA - PINE ISLAND | 81 | 82 |
|        | 2004 Average     | 86 | 89 |

| Grand Mean | 85 | 87 |
|------------|----|----|
|            |    |    |

Table 6. Seedling vigor for CL131 and CL161 across several trials at multiple locations in Louisiana (2003-2004).

|      | CL - LAKE ARTHUR  2003 Average | 8    | 4     |
|------|--------------------------------|------|-------|
| 2003 |                                | 6    | 4     |
| YEAR | TEST                           | CEST | ©L161 |

|      | CL - JEFF DAVIS  | 4 | 4 |
|------|------------------|---|---|
|      | CL - LAKE ARTHUR | 5 | 5 |
| 2004 | CL - PINE ISLAND | 5 | 4 |
|      | CA - LAKE ARTHUR | 4 | 4 |
|      | CA - PINE ISLAND | 3 | 3 |
| :    | 2004 Average     | 5 | 4 |

| 0          |    | 07 |
|------------|----|----|
| Grand Mean | 86 | 87 |

Table 7. Reaction of CL131 and CL161 to Sheath Blight (*Rhizoctonia solani*) in field plots (2004).

| YEAR TEST     | CL131 | e e c |
|---------------|-------|-------|
| 2004 CA - RRS | 7.5   | 7.3   |

<sup>\*</sup> Using a scale of 0 = very resistant to 9 = very susceptible.

Table 8. Reaction of CL131 and CL161 to Leaf Blast (*Pyricularia grisea*) in field plots (2004).

| YEAR TEST     | e-131 |     |
|---------------|-------|-----|
| 2004 CA - RRS | 1.8   | 4.5 |

<sup>\*</sup> Using a scale of 0 = very resistant to 9 = very susceptible.

Table 9. Reaction of CL131 and CL161 to the physiological disorder straighthead (2004).

| YEAR TEST | CL131 | CL161 |
|-----------|-------|-------|
| 2004 RRS  | 5.7   | 3.0   |

Table. 10 Rough, brown and milled grain dimensions and weight of CL131 and CL161 grown in Crowley, LA (2004).

| Variety | Туре   | Length<br>mm | Width<br>mm | Thickness<br>mm | L/W Ratio |
|---------|--------|--------------|-------------|-----------------|-----------|
|         | Rough  | 9.06         | 2.53        | 2.03            | 3.58      |
| CL131   | Brown  | 7.14         | 2.22        | 1.72            | 3.22      |
|         | Milled | 6.62         | 2.18        | 1.67            | 3.04      |
|         |        |              |             |                 |           |
|         | Rough  | 9.10         | 2.52        | 2.00            | 3.61      |
| CL161   | Brown  | 6.96         | 2.17        | 1.69            | 3.21      |
|         | Milled | 6.65         | 2.07        | 1.64            | 3.21      |

Table 11. Reaction of CL131 and CL161 to Narrow Brown Leaf Spot\* (*Cercospora janseana*) in field plots (2004).

| YEAR TEST     | CL131 | CL161 |
|---------------|-------|-------|
| 2004 CA - RRS | 5.0   | 3.5   |

<sup>\*</sup> Using a scale of 0 = very resistant to 9 = very susceptible.

Table 12. Reaction of CL131 and CL161 to Leaf Smut\* (*Entyloma oryzae*) in field plots (2004).

| YEAR TEST     | CL131 | CL161 |
|---------------|-------|-------|
| 2004 CA - RRS | 5.8   | 3.8   |

<sup>\*</sup> Using a scale of 0 = very resistant to 9 = very susceptible.

Table 13. Reaction of CL131 and CL161 to Brown Spot\* (*Cochiobolus miyabeanus*) in field plots (2004).

| YEAR TEST     | CL131 | CHA |
|---------------|-------|-----|
| 2004 CA - RRS | 1.5   | 1.8 |

<sup>\*</sup> Using a scale of 0 = very resistant to 9 = very susceptible.

| Trait            | Perfor | mance | Number<br>of Tests | Reference |
|------------------|--------|-------|--------------------|-----------|
|                  |        | •     | 7                  |           |
|                  | CL131  | CL161 |                    |           |
| Yield            | 6520   | 6455  | 15                 | Table 1   |
| Ratoon           |        |       |                    |           |
|                  |        |       |                    |           |
| Whole            | 67.2   | 66.2  | 9                  | Table 2   |
| Total            | 72.1   | 70.8  | 9                  | Table 3   |
|                  |        |       |                    |           |
| Length-Rough     | 9.06   | 9.10  |                    | Table 10  |
| Width-Rough      | 2.53   | 2.52  |                    | Table 10  |
| L/W Ratio-Rough  | 3.58   | 3.61  |                    | Table 10  |
|                  |        |       |                    |           |
| Length-Brown     | 7.14   | 6.69  |                    | Table 10  |
| Width-Brown      | 2.22   | 2.17  |                    | Table 10  |
| L/W Ratio-Brown  | 3.22   | 3.21  |                    | Table 10  |
|                  |        |       |                    |           |
| Length-Milled    | 6.62   | 6.65  |                    | Table 10  |
| Width-Milled     | 2.18   | 2.07  |                    | Table 10  |
| L/W Ratio-Milled | 3.04   | 3.21  |                    | Table 10  |
|                  |        |       |                    |           |
| Vigor            | 5      | 4     | 7                  | Table 6   |
| Height (cm)      | 90     | 103   | 14                 | Table 4   |
| Days to 50%      | 85     | 87    | 15                 | Table 5   |
|                  |        |       |                    |           |
| Blast (LB/RNB)   | 1.8    | 4.5   |                    | Table 8   |
| Sheath Blight    | 7.5    | 7.3   |                    | Table 7   |
| Straighthead     | 5.7    | 3.0   |                    | Table 9   |

| REPRODUCE LOCALLY. Include form number and edition date on all  | reproductions F   | ORM APPROVED - OMB No. 0581-005                                  |
|---|---|--|
| U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE  EXHIBIT E  | Application is required in order to detectificate is to be issued (7 U.S.C. 24 confidential until the certificate is issued | (21). The information is held                                    |
| STATEMENT OF THE BASIS OF OWNERSHIP   |   |  |
| 1. NAME OF APPLICANT(S)   | 2. TEMPORARY DESIGNATION<br>OR EXPERIMENTAL NUMBER  | 3. VARIETY NAME  |
| Louisiana State University Agricultural Center  | CL020 CL131   |  |
| 4. ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP, and Country)   | 5. TELEPHONE (Include area code) 6. FAX (Include area code)   |  |
| Rice Research Station<br>1373 Caffey Road   | (337) 788-7531  | (337) 788-7553   |
| Rayne, LA 70578   | 7. PVPO NUMBER 0050   | 0349   |
| 8. Does the applicant own all rights to the variety? Mark an "X" in the   | appropriate block. If no, please explai   | n. YES NO  |
| 9. Is the applicant (individual or company) a U.S. national or a U.S. ba  | ased company? If no, give name of co  | ountry. YES NO   |
| 10. Is the applicant the original owner? YES  | NO If no, please answer <u>one</u> o  | of the following:  |
| a. If the original rights to variety were owned by individual(s), is (a   | NO If no, give name of count  |  |
| b. If the original rights to variety were owned by a company(ies),  | is (are) the original owner(s) a U.S. bas   |  |
| 11. Additional explanation on ownership (Trace ownership from original  | al breeder to current owner. Use the re   | verse for extra space if needed):                                |
|   |   |  |
|   |   |  |
| •   |   |  |
|   |   |  |
| PLEASE NOTE:  |   |  |
| Plant variety protection can only be afforded to the owners (not license  | es) who meet the following criteria:  |  |
| If the rights to the variety are owned by the original breeder, that per<br>national of a country which affords similar protection to nationals of                                  | rson must be a U.S. national, national o  | f a UPOV member country, or s.                                   |
| <ol><li>If the rights to the variety are owned by the company which employe<br/>nationals of a UPOV member country, or owned by nationals of a co-<br/>genus and species.</li></ol> | ed the original breeder(s), the company<br>ountry which affords similar protection to                                       | must be U.S. based, owned by onationals of the U.S. for the same |
| 3. If the applicant is an owner who is not the original owner, both the o   | riginal owner and the applicant must me   | eet one of the above criteria.                                   |
| The original breeder/owner may be the individual or company who dire<br>Act for definitions.  | cted the final breeding. See Section 4  | (a)(2) of the Plant Variety Protection                           |
| According to the Paperwork Reduction Act of 1995, an agency may not conduct or soonsor a  | nd a person is not required to respond to a collection  | onf information unless it displays a valid OMP                   |

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information collection is 0581-0055. The time required to complete this information collection is estimated to average 0.1 hour per response, including the time for reviewing the instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, sexual orientation, marital or family status, political beliefs, parental status, or protected genetic information. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, D.C. 20250-9410 or call (202) 720-5964 (voice and TDD). USDA is an equal opportunity provide and employer.